

*A review of the important adverse biological effects of atmospheric photochemical smog on man and animals expressed in terms of atmospheric "total oxidant" concentrations is presented, based on published reports and some yet unpublished reports and research findings. Additionally, data based on laboratory exposures of man and animals to the pure agents, ozone and peroxyacyl nitrates (PAN compounds), important components of the "total oxidant" mixture which have been identified in photochemical smog, are reviewed.*

## **THE BIOLOGICAL EFFECTS OF PHOTOCHEMICAL AIR POLLUTANTS ON MAN AND ANIMALS**

*Louis S. Jaffe, M.A., F.A.P.H.A.*

**A**IR pollution, the unfavorable alteration of one of our most important natural resources—the air we breathe—consists of emissions of gaseous and particulate products often followed by secondary reactions in the atmosphere. Except for natural causes, polluted air is a result of man's activities: industrialization, urbanization, transportation, heating of homes and buildings, and burning of wastes.

While the increasing pollution of our limited air supply has become a matter of national concern, the problem is far from new. For example, an important type of air pollution has existed and been a problem for hundreds of years. From the beginning of the 14th century to the early part of the 20th century, air pollution by coal smoke and gases has been the predominant form, and in many industrialized communities and areas of the world still is the principal form of air pollution.<sup>1</sup> This type of air pollution, where smoke, particulates of various sizes, and

gases, primarily sulfur oxides, are produced from the combustion of coal or other fossil fuels, is characterized by its chemically reducing action and is often identified as "London-type smog." The "London-type smog" is the dominant kind of atmospheric pollution found in such large urban industrialized communities as New York, Philadelphia, and Chicago, and foreign urban communities where coal and other fossil fuels are consumed.

Another major type of community air pollution is photochemical air pollution, often called photochemical smog. This type of pollution is formed as a result of atmospheric reactions initiated by ultraviolet radiation from sunlight on certain gaseous emissions, the oxides of nitrogen and reactive organic compounds, derived from the combustion of organic fuels. The motor vehicle is a principal source of these emissions.<sup>2,24</sup> Other sources are fuel burning for heat and power, refuse burning, evaporation losses

from petroleum products, and the industrial and commercial use of organic solvents.<sup>2</sup>

Photochemical air pollution is of recent origin, historically speaking. The first manifestations of this form of community air pollution: excessive cracking of rubber products,<sup>4,5,10,15</sup> a peculiar form of leaf injury to many agricultural and ornamental crops,<sup>3,5-11</sup> and eye irritation,<sup>5,10-12,15</sup> were noted in the Los Angeles area only a little more than 20 years ago, hence the commonly used designation, "Los Angeles-type smog." Other expressions of photochemical smog are respiratory distress,<sup>13,14,54</sup> haze formation causing a reduction in visibility with consequent hazards to ground, air, and water transportation,<sup>10-12,15-17,24</sup> and a characteristic objectionable odor,<sup>17</sup> as well as the presence of unusually high levels of atmospheric photochemical oxidants. More recent findings of effects on man and animals will be described in some detail below.

The photochemical oxidants are a major class of gaseous compounds found in photochemical smog. They consist of a complex mixture of atmospheric oxidizing substances which vary in composition with time and place; all the components are not yet completely defined chemically. Nonetheless, they can, with appropriate analytical procedures, be measured routinely and analyzed collectively as "total oxidant," which serves as a useful index of photochemical air pollution. The rate of formation of photochemical oxidants is controlled by the intensity of sunlight, and the concentrations of the major reactant gases—the reactive organic compounds (olefins, aromatic hydrocarbons, and aldehydes) and nitrogen oxides.<sup>15,19,20</sup> The accumulation of these reactant gases, in turn, is determined by their emission rates from the several sources, as well as by meteorological and topographical factors. Meteorological factors, such as low wind speed and vertical stability, play an important

part in the accumulation of photochemical air pollutants in the community air.<sup>21</sup> Temperature plays a significant but secondary role in the rate of accumulation of photochemical air pollutants: all chemical reactions are accelerated with increased temperature.<sup>18,19</sup>

The adverse effects of photochemical smog now occur in every region of the United States. Higher community photochemical oxidant concentrations and greater frequency of other photochemical smog manifestations, however, in general are found more frequently in the lower latitudes where ample sunshine and higher temperatures prevail year round.<sup>21</sup>

An "oxidant," as used in the air pollution field, may be defined as a substance which oxidizes a selected reagent under specified conditions. Currently, the field methods commonly used for routine measuring of the ambient photochemical oxidants in the community atmosphere measure "total oxidant," i.e., the collective net oxidizing effect of all the oxidizing substances formed in the atmosphere, indiscriminately. Oxidant values, measured as "total oxidant" usually by the neutral potassium iodide method, serve as practical indexes of the various biological and material expressions of photochemical smog, i.e., when the "total oxidant" level is high the various signs and symptoms associated with photochemical smog are more pronounced; when the "total oxidant" level is low, the associated manifestations of photochemical smog are minimal.

Using more sophisticated instrumentation, however, some of the individual atmospheric photochemical oxidants have been determined. Ozone has been identified as a principal oxidant produced in photochemical smog.<sup>4,5,15,17-20,22,23</sup> Other atmospheric oxidants which have been isolated and identified are peroxyacyl nitrates, a related group of organic peroxidic nitrogen compounds,<sup>25-28</sup> designated as PAN compounds, or more recently as PaNs.<sup>29</sup> The PaNs have been determined

by Stephens and his associates to be a homologous series of compounds; they have identified peroxyacetyl nitrate (PAN), the principal compound in the group, as well as peroxypropionyl nitrate (PPN), and peroxybutyryl nitrate (PBN), thus far.<sup>25-28</sup> Other oxidants present in photochemical smog are nitrogen oxides, which act as weak oxidants, other organic peroxides, free radicals, and other compounds yet unidentified.

Daily fluctuation of "total oxidant" is a major characteristic of photochemical smog and follows a distinctive pattern: low concentrations at night are followed by an increase in the early morning; a peak reached during the remainder of the sunlit hours, usually in the afternoon, is followed by slow decay and a gradual disappearance after sunset.<sup>12,17,30,31</sup> At peak levels of "total oxidant," ozone may constitute up to 90 per cent or more of this total (neutral KI methods).<sup>32,33</sup> An annual cycle exists; high hourly concentrations are found more frequently, usually from May through October, than during the rest of the year.<sup>30,34</sup> Peak "total oxidant" levels up to about 1.0 ppm in dense community photochemical smog have been reported.<sup>35</sup>

Some of the more recent studies and findings on the effects of atmospheric photochemical oxidants expressed in terms of "total oxidant" levels (neutral KI method) will be briefly described. Data on the effects of ozone and PAN based on laboratory or chamber exposures of humans and of animals to the generated pure agents are also included because of their identification as important oxidants found in photochemical smog. The data based on exposures to the single oxidizing agents in pure form are supplementary to the data based on ambient air studies expressed as "total oxidant." An understanding of their effects contributes substantially to our knowledge of the effects of the "total oxidant" mixture.

Substantial data on the adverse effects

of generated ozone on man and animals are available. Ozone has been recognized as an important industrial toxicant for a long time, much before an awareness of its presence as an important component in photochemical smog was identified. It is formed in nature by high voltage discharge during thunderstorms and is formed in natural processes by the photodissociative action of solar ultraviolet short wavelength radiation on the oxygen molecules in the lower stratosphere.<sup>36,37</sup> It has been encountered occupationally in appreciable levels in inert gas shielded arc-welding devices<sup>38-42</sup>; and it is formed by high voltage discharge from the brushes of motors, neon signs, and other electrical equipment. It has been used in the past and to a limited extent at present for air purification and odor control in buildings and homes by generation from ozonizing equipment<sup>43-47</sup>; in cold storage plants for the control of molds and bacteria<sup>48</sup>; and in the treatment of sewage wastes,<sup>49</sup> and purification of drinking water.<sup>50</sup> On the other hand, the PANs have neither been found in nature nor been identified as contaminants in other areas of environmental health excepting air pollution. In addition, their synthesis and generation for purposes of laboratory studies and their methods of measurement are much more difficult than those for ozone; hence, comparatively little is known about them at present.

## Effects of Photochemical Smog on Man

### *Short-Term Effects*

#### *Eye Irritation*

The most commonly experienced and recognized attribute of photochemical smog by man is eye irritation. The components causing eye irritation have not been completely identified as yet but there is a significant correlation between eye irritation and levels of atmospheric "total oxidant" based on eye irritation surveys.<sup>75,76</sup> Eye irritation is experienced

at community oxidant levels of 0.10 ppm and higher by substantial elements of the population.<sup>51</sup> To a high degree, eye irritation is subjective; it is influenced by abundance of light, humidity, haze, and wind currents, as well as the somatic condition of the observer.<sup>24</sup>

#### **Odor**

There is a characteristic pungent odor associated with photochemical smog.<sup>5,52</sup> Ozone is an acrid component of this odor.<sup>17,52</sup>

#### *Prolonged Effects*

##### **Respiratory Effects**

Recent studies have shown that it is harder for humans, especially patients suffering from chronic respiratory disease, to breathe in areas having even a moderate level of photochemical air pollution.<sup>13,54,78</sup> Patients with chronic bronchopulmonary disease, when exposed in clinical studies to smoggy Los Angeles air having 0.138 ppm "total oxidant" (average of the daily maxima) for one week, showed a significant and uniform increase in oxygen consumption and a decrease in arterial blood oxygen tension (oxygen partial pressure) during light exercise compared with the findings in each patient while breathing clean filtered air for a similar period.<sup>13,54</sup> Thus, although the patients were consuming more oxygen, less of it was being made available to the body. These patients also had greater difficulty in breathing (increased pulmonary airway resistance and work of breathing) when breathing the smoggy air.<sup>13,54</sup>

#### **Effects of Photochemical Smog on Animals**

##### *Short-Term Effects*

Studies on guinea pigs exposed over their lifetime to atmospheric photochemi-

cal smog showed that the animals experienced greater difficulty in breathing on days of high photochemical air pollution levels (0.30 ppm "total oxidant" and above for three hours).<sup>55</sup> Even more significant increases in breathing resistance, particularly in the older animals, occurred on days having severe smog (0.50 ppm peak "total oxidant" levels and above,<sup>56</sup> the first alert level in Los Angeles).<sup>11</sup> Temporary ultrastructural alterations in the wall cells of lung alveoli of nine-month-old mice occurred after exposure to a severe smog episode (>0.40 ppm "total oxidant" for 2-3 hours).<sup>79</sup> Recovery of the alveolar tissues took place within 14 hours.<sup>79</sup> Laboratory mice exposed to a diluted irradiated auto exhaust atmosphere containing as low as 0.15 ppm mean "total oxidant" for four hours and then to an infectious *Streptococcus aerosol* showed a significant increase (enhancement) of mortality (27 per cent) over similarly infected control animals exposed first to filtered air.<sup>77</sup>

##### *Prolonged Effects*

Laboratory mice (A and A/J strains) exposed to community photochemical smog levels (0.14 ppm average of the daily "total oxidant" maxima) over a 16-month period showed an increase in lung tumor development (adenoma) in the aging animals when compared to controls exposed to filtered air.<sup>57</sup>

Several strains of young laboratory mice, five to eight weeks old, exposed continuously to low levels of synthetic smog (0.08 ppm average of the "total oxidant" daily maxima, irradiated auto exhaust cycled within test chambers to simulate the daily oxidant fluctuation pattern occurring in the atmosphere) showed increased susceptibility to pulmonary infection and chronic disease during the latter half of the animals' lifetime compared to controls exposed only to clean filtered air.<sup>58</sup> Significant de-

creases in mouse fertility and survival rate of infant LAF<sub>1</sub> strain mice were also manifested during the 13-month exposure period (through the animals' effective reproductive period).<sup>58</sup>

## Effects of Ozone on Man

### *Short-Term Effects*

#### Odor

The characteristic sharp odor of ozone, sometimes described as an "electrical" odor, can be detected instantaneously at very low concentrations ( $<0.02$ - $0.05$  ppm), depending on individual acuity.<sup>53,59-60,44</sup> At somewhat higher concentrations ( $0.05$ - $0.10$  ppm) the odor becomes more pronounced and disagreeable.<sup>53,60</sup>

#### Respiratory Effects

Ozone is a severe irritant to all mucous membranes. Significant exposures may cause pulmonary congestion and other complications. The first symptoms of irritation due to ozone, dryness of the upper respiratory passages and initial irritation to the mucous membranes of the nose and throat occur after brief exposures (13-30 minutes) to low concentrations ( $0.05$ - $0.10$  ppm).<sup>53,44</sup> At higher concentrations ( $0.30$ - $1.0$  ppm) all within the range of community oxidant levels and exposures for longer intervals (15 minutes-2 hours), marked respiratory irritation is accompanied by respiratory distress (choking, coughing and severe fatigue) particularly so at the upper end of this range.<sup>53,61-63</sup>

At relatively high concentrations, equivalent to the ozone content in severe photochemical smog, lung function is impaired for the duration of exposure and for some time after. Normal subjects exposed to a range of  $0.6$ - $0.8$  ppm ozone in single two-hour periods showed a marked change in lung function (highly significant reduction in diffusing capacity (25 per cent) and significant re-

duction in vital capacity and forced expiratory volume) lasting up to 24 hours.<sup>41</sup> The subjects developed bronchial irritation and substernal soreness lasting up to 12 hours, as well as a slight dry cough disappearing within 24 hours. At still higher concentrations, beyond the maximum levels presently found in community photochemical smog ( $1.5$ - $2.0$  ppm), a single two-hour exposure to ozone caused general morbidity in a man lasting for approximately two weeks.<sup>64</sup> His immediate symptoms included impaired lung function, severe chest pains, altered taste sensation, coughing, headache, and extreme fatigue. During the period of severe malaise following exposure, the subject also showed a loss of coordinating ability and difficulty in expression and articulation.<sup>64</sup>

#### Effects on Vision

Humans when exposed to ozone concentrations as low as  $0.20$  ppm ( $0.20$ - $0.50$  ppm) for three hours showed a considerable decrease in visual acuity and changes in the extra-ocular muscle balance, reduced night vision, lateral phoria (divergence), and other changes in visual parameters.<sup>65</sup> These effects were all manifested at concentrations of  $0.20$  in three hours; effects were more pronounced at  $0.35$  and  $0.50$  ppm respectively.

### *Prolonged Effects*

Few prolonged ozone exposure studies of humans have been reported. Occupational exposures of welders to intermittent relatively high concentrations of ozone (range of  $0.3$ - $1.7$  ppm) over a two-week period caused severe recurrent headache, fatigue, chest pains, difficulty in breathing and wheezing.<sup>40,66</sup>

## Effects of Ozone on Animals

Ozone is a strong irritant of the mucous membranes of animals. It augments

the morbidity effects of respiratory infection (and shortens the morbidity span) in test animals exposed both to the respiratory infection and to ozone. Recent findings indicate that ozone also may produce secondary systemic effects on body metabolism and function, including the inhibition of normal thyroid function.<sup>67</sup>

### *Short-Term Effects*

A mean ozone concentration as low as 0.08 ppm for three hours caused a significant increase in mortality of Streptococcus-infected test animals over similarly infected animals not exposed to ozone.<sup>68</sup> In a recent study on the enhancement of respiratory disease due to prior exposure to ozone, Coffin exposed groups of Swiss mice each weighing 25 g. to a specific level of ozone for three hours followed by exposure for half an hour to an aerosolized culture of Streptococcus in a series of ozone levels ranging from 0.05 to 0.50 ppm. The control animals were exposed for three hours to filtered air and then to the Streptococcus aerosol for half an hour. Mortality was measured over a ten-day period. The increase in mortality after exposure to 0.08 ppm ozone for three hours and then the Streptococcus aerosol was 23 per cent over the mortality of the controls. (P value <0.05.)

Concentrations of ozone comparable to commonly found atmospheric community pollution levels also caused a decreased activity in mice (decreased running ability after exposure to 0.20 ppm for 6 hours), while ozone levels of 0.34 ppm and above for two hours caused temporarily impaired lung function.<sup>69</sup> At ozone concentrations well above present community levels (1.50 ppm and above), pathological lung changes occurred in previously healthy mice after a single four-hour exposure which apparently were reversible after withdrawal from the ozone environment.<sup>70</sup>

### *Prolonged Effects*

Continuous ozone exposure for 3-17 weeks at concentrations of 0.10-0.25 ppm shortened the lives of infected guinea pigs and increased their mortality rate,<sup>71</sup> while intermittent prolonged exposures (0.10-0.20 ppm for seven hours/day, five days/week for three weeks) caused an increase in the mortality of newborn mice.<sup>72</sup> Significant increased mortality and severe chronic lung injury including hemorrhage, fibrosis of the lung parenchyma and constriction of the lung airways occurred in guinea pigs and rats when intermittently exposed to ozone concentrations of 1.50 ppm (neutral KI method) for 62 weeks.<sup>73</sup>

### **Effects of PAN (Peroxyacetyl Nitrate) and PAN Compounds (Peroxyacetyl Nitrates) on Man and Animals**

Since the PAN compounds have been used in human or animal exposures for only a short time, comparatively little information has been developed to date.

### **Effects of PAN or PAN Compounds on Man**

Peroxyacetyl nitrate (PAN) appears to have an effect on pulmonary function of humans similar to that reported for ambient oxidant. In a recent study, healthy students when performing moderate exercise and breathing 0.30 ppm PAN for five minutes showed a statistically significant increase in oxygen uptake over the oxygen consumption of the same students during an identical period when breathing clean filtered air during exercise; breathing (expiration velocity during the recovery stage) was also affected.<sup>74</sup>

PAN has been reported to cause eye irritation when panels of human volunteers were exposed to pure synthetic PAN at a concentration of 1.0 ppm for 10-15 minutes; the threshold level of

detection of synthetic PAN as an eye irritant was 0.5 ppm for 12 minutes.<sup>26</sup> For some reason not completely clear, the levels of PAN and other known or suspected lachrymatory agents found in photochemical smog, when tested in pure form, require a higher concentration to produce eye irritation than the concentrations of these reactants found in ambient photochemical smog.<sup>15,75</sup>

### Effects of PAN on Animals

No apparent evidence of pronounced injury to animals exposed to PAN in concentrations approximating atmospheric levels has been observed to date. These studies are continuing.

### Conclusion

When concentrations of photochemical oxidants are used as measures of photochemical smog intensity, a number of specific biological effects on man and animals based on both atmospheric and laboratory studies have been documented. Photochemical oxidants, such as frequently found in urban community atmospheres as measured by continuous air monitoring instruments operated by federal, state, and local agencies, cause a repeated and continuing biological impact on man and animals of varying degree and frequency in every region of the country.

### REFERENCES

- Chambers, L. A. "Classification and Extent of Air Pollution Problems." In: *Air Pollution*, Vol. I, pp. 4-22. Stern, A. C., editor. New York, N. Y.: Academic Press, 1962.
- Leighton, P. A. Man and Air in California. Proceedings, Conference on Man in California, 1980's. Jan. 27, 1964. Sacramento, Calif. University of California, Extension Division, Berkeley.
- Middleton, J. T.; Kendrick, J. B.; and Schwalm, H. W. Injury to Herbaceous Plants by Smog or Air Pollution. US Department of Agriculture Plant Disease Reporter. 34:245, 1950.
- Bradley, C. E., and Haagen-Smit, A. J. The Application of Rubber in the Quantitative Determination of Ozone. J. Rubber Chem. Tech. 24:750, 1951.
- Haagen-Smit, A. J. Chemistry and Physiology of Los Angeles Smog. Ind. Eng. Chem. 44:61342 (June), 1952.
- Haagen-Smit, A. J.; Darley, E. F.; Zaitlin, M.; Hull, H.; and Noble, W. Investigations on Injury to Plants from Air Pollution in the Los Angeles Area. Plant Physiol. 27:18, 1952.
- Bobrov, R. A. Effect of Smog on the Anatomy of Oak Leaves. Phytopath. 42:558, 1952.
- Middleton, J. T.; Kendrick, J. B.; and Darley, E. F. Air Pollution Injury to Crops. California Agric. 7,11, 1953.
- \_\_\_\_\_. Air-borne Oxidants as Plant Damaging Agents. In: Proceedings of the Third National Air Pollution Symposium. Stanford Research Institute, Pasadena, Calif., 1955, pp. 191-197.
- Los Angeles County Air Pollution Control District: Second Technical and Administrative Report on Air Pollution Control in Los Angeles County, 1950-1951. Los Angeles, Calif., 1952.
- State of California Department of Health Technical Report. California Standards for Ambient Air Quality and Motor Vehicle Exhaust. Berkeley, Calif., 1960.
- Dickinson, J. E. Air Quality of Los Angeles County. Technical Progress Report II. Los Angeles County Air Pollution Control District (Feb.), 1961.
- Remmers, J. E., and Balchum, O. J. Effects of Los Angeles Urban Pollution Upon Respiratory Function of Emphysematous Patients. Paper 65-43. Presented at the 58th annual meeting, Air Pollution Control Association, Toronto, Canada, June, 1965.
- Motley, H. L.; Smart, R. H.; and Leftwich, C. I. Effect of Polluted Los Angeles Air (Smog) on Lung Volume Measurements. J.A.M.A. 171:1469, 1959.
- Wayne, L. G. The Chemistry of Urban Atmospheres, Technical Progress Report III. Los Angeles Air Pollution Control District (Dec.), 1962.
- Mader, P. P.; MacPhee, R. D.; Lofberg, R. T.; and Larson, G. P. Composition of Organic Portions of Atmospheric Aerosols in the Los Angeles Area. Ind. Eng. Chem. 44:1352 (June), 1952.
- Haagen-Smit, A. J. Photochemistry and Smog. J. Air Pollut. Contr. Ass. 13:9:444 (Sept.), 1963.
- Leighton, P. A. Physical Chemistry, Vol. IX. Photochemical Aspects of Air Pollution. New York, N. Y.: Academic Press, 1961.
- Altschuler, A. P., and Bufalini, J. J. Photochemical Aspects of Air Pollution: A Review. Photochem. and Photobiol. 4:97 (Jan.), 1965.
- Haagen-Smit, A. J.; Bradley, C. E.; and Fox, M. M. Ozone Formation in Photochemical Oxidation of Organic Substances. Ind. Eng. Chem. 45:2086 (Sept.), 1963.
- Neiburger, M. Meteorological Aspects of Air Pollution: Their Relationship to Biological Responses. Arch. Environ. Health 14:1:41 (Jan.), 1967.
- Haagen-Smith, A. J., and Fox, M. M. Ozone Formation in Photochemical Oxidation of Organic Substances. Ind. Eng. Chem. 48:1484 (Sept.), 1956.
- Richards, L. M. The Formation of Ozone in Polluted Atmospheres. J. Air Pollut. Contr. Ass. 5:4:215 (Feb.), 1956.
- Haagen-Smit, A. J. Atmospheric Reactions of Air Pollutants. Ind. Eng. Chem. 48:65A (Dec.), 1956.
- Darley, E. F.; Stephens, E. R.; Middleton, J. T.; and Hanst, P. L. Oxidant Plant Damage from Ozone-Olefin Reactions. Int. J. Air Poll. 1:155, 1959.
- Stephens, E. R.; Darley, E. F.; Taylor, O. C.; and Scott, W. E. Photochemical Reaction Products in Air Pollution. Ibid. 4, 1-2, 79, 1961.
- Darley, E. F.; Ketner, K. A.; and Stephens, E. R. Analyses of Peroxyacyl Nitrates by Gas Chromatography with Electron Capture Detection. Anal. Chem. 35:589 (Apr.), 1963.
- Stephens, E. R. Absorptivities for Infrared Determination of Peroxyacyl Nitrates. Ibid. 36:928 (Apr.), 1964.
- Jaffe, L. S. Effects of Photochemical Air Pollution on Vegetation with relation to Air Quality Requirements. J. Air Pollut. Contr. Assoc. 17:1:38 (Jan.), 1967.
- Cholak, J.; Schafer, L. J.; Younker, W. J.; and Yeager, D. W. Pollution in a Number of American

- Cities. Arch. Environ. Health 11:280 (Apr.), 1955.
31. Stephens, E. R. Temperature Inversions and the Trappings of Air Pollutants. Weatherwise (Am. Meteorological Society) 18.4:172 (Aug.), 1965.
32. Renzetti, N. A., and Romanovsky, J. C. A Comparative Study of Oxidants and Ozone in the Los Angeles Atmosphere. Arch. Environ. Health 14:458, 1956.
33. Cherniack, I., and Bryan, R. J. A Comparison of Various Types of Ozone and Oxidant Detectors which are used for Atmospheric Air Sampling. J. Air Pollut. Contr. Assoc. 15:8:351 (Aug.), 1965.
34. Hamming, W. J.; MacPhee, R. D.; and Taylor, J. R. Contaminant Concentrations in the Atmosphere of Los Angeles County. Ibid. 10:7 (Feb.), 1960.
35. Tebbens, B. D. "Residual Pollution Products in the Atmosphere." In: Air Pollution, Vol. I, p. 35. A. C. Stern, editor. New York, N. Y.: Academic Press, 1962.
36. Jaffe, L. S., and Estes, H. D. Ozone in High-Altitude Aircraft Cabins. Arch. Environ. Health 9:61 (July), 1964.
37. Regener, V. H. Vertical Distribution of Atmospheric Ozone. Nature 167:276, 1951.
38. Ferry, J. J., and Ginther, G. B. Gases Produced by Inert-Arc-Welding. Am. Indust. Hyg. A. Quart. 13: 196, 1952.
39. Kleinfeld, M., and Giel, C. P. Clinical Manifestations of Ozone Poisoning. Report of a New Source of Exposure. Am. J. M. Sc. 23:638, 1956.
40. Challen, P. J. R.; Hickish, D. E.; and Bedford, J. An Investigation of Some Health Hazards in an Inert-Gas, Tungsten Arc Welding Shop. Brit. J. Indust. Med. 15:276, 1958.
41. Young, W. A.; Shaw, D. B.; and Bates, D. V. Effects of Low Concentrations of Ozone on Pulmonary Function. J. Appl. Physiol. 19:765, 1964.
42. Frant, R. Formation of Ozone in Gas-Shielded Welding. Ann. Occup. Hyg. 6:113, 1963.
43. Stokinger, H. E. Ozone Toxicology; A Review of Research and Industrial Experience: 1954-1964. Arch. Environ. Health 10.5:719 (May), 1965.
44. Witheridge, W. W., and Yaglou, C. P. Ozone in Ventilation—Its Possibilities and Limitations. J. Am. Soc. Heat. & Vent. Engin. 45:509, 1939.
45. Jordan, E. O., and Carlson, A. J. Ozone: Its Bactericidal, Physiologic and Deodorizing Action. J.A.M.A. 61:1007 (Sept. 27), 1913.
46. Elford, W. J., and van den Ende, J. Investigation on the Merits of Ozone as an Aerial Disinfectant. J. Hyg. 42:240, 1942.
47. Ebermeyer, M. De l'ozone atmospherique (Atmospheric Ozone). Le Moniteur Scientifique (3rd series) 15:891, 1873, Paris, France.
48. Nagy, R. "Application of Ozone from Sterilamp in Control of Molds, Bacteria and Odors." In: Ozone Chemistry and Technology. Advances in Chemistry Series. 21:57 (Mar.), 1959. American Chemical Society. Washington, D. C.
49. Guinvarc'h, P. "Three Years of Ozone Sterilization of Water in Paris." Ibid.
50. Torricelli, A. "Drinking Water Purification." Ibid.
51. Faith, W. L.; Renzetti, N. A.; and Rogers, L. H. Third Technical Progress Report. Air Pollution Foundation Report No. 17, Los Angeles, Calif. (Mar.), 1957.
52. Haagen-Smit, A. J. "Reactions in the Atmosphere." In: Air Pollution, Vol. I, pp. 41-46. New York, N. Y.: Academic Press, 1962.
53. Henschler, A.; Stier, H.; Beck, H.; and Newmann, W. Olfactory Threshold of Some Important Irritant Gases and Manifestations in Man by Low Concentrations. Archiv. fur Gewerbepathologie und Gewerbehygiene. 17.6:547 (Mar.), 1960.
54. Balchum, O. J. Personal communication, May 19, 1966. Data to be published in Arch. Environ. Health. In press.
55. Swann, H. E., and Balchum, O. J. The Biological Effects of Urban Air Pollution. IV. Effects of Acute Smog on Respiration of Guinea Pigs. Arch. Environ. Health 12:698 (June), 1966.
56. Swann, H. E.; Brunol, D.; Wayne, L. G.; and Balchum, O. J. Biological Effects of Air Pollution. II. Chronic Exposure of Guinea Pigs. Ibid. 11.6:765 (Dec.), 1965.
57. Gardner, M. B. Biological Effects of Urban Air Pollution. III. Lung Tumors in Mice. Ibid. 12:305 (Mar.), 1966.
58. Hueter, F. G.; Contner, G. L.; Busch, K. A.; and Hinners, R. G. Biological Effects of Atmospheres Contaminated by Auto Exhaust. Ibid. 12:553 (May), 1966.
59. Stokinger, H. E. Ozone Toxicity. A Review of the Literature Through 1953. Arch. Indust. Hyg. Occ. Med. 9:366, 1954.
60. Scott, D. B. M., and Leshner, E. C. Effect of Ozone on Survival and Permeability of Escherichia coli. J. Bact. 85:567, 1963.
61. Dadlez, M. J. Ozone Formed by Ultra-Violet Rays. Quart. J. Pharmacy. 1:99, 1928 (Also: Ozone dans l'air. Recherche sur sa production par les rayons ultraviolets. L'Union Pharmaceutique 69:17, 1928).
62. Flury, F., and Zernik, F. Schädliche Gase, Dampfe, Nebel, Rauch und Staubarten. (Noxious Gases, Fumes, Vapors, and Varieties of Smoke and Dust.) Berlin, Germany: Springer, 1931.
63. Wilksa, S. Ozone. Its Physiological Effects and Analytical Determination in Laboratory Air. Acta chem. scandinav. 1:359, 1951.
64. Griewold, S. S.; Chambers, L. A.; and Motley, H. L. Report of a Case of Exposure to High Ozone Concentrations for Two Hours. Arch. Indust. H. 15:108, 1957.
65. Lagerwerff, J. M. Prolonged Ozone Inhalation and Its Effects on Visual Parameters. Aerospace Medicine 34.6:479 (June), 1963.
66. Kleinfeld, M.; Giel, C.; and Tabershaw, I. R. Health Hazards Associated with Inert-Gas-Shielded Metal Arc Welding. Arch. Indust. H. 15:27, 1957.
67. Fairchild, E. J. II: Tolerance Mechanisms: Determinants of Lung Response to Injurious Agents. Arch. Environ. Health 14.1:111 (Jan.), 1967.
68. Coffin, D. L. Effect of Ozone on Enhancement of Bacteriological Infection. Ibid. In press, 1967.
69. Murphy, S. D.; Ulrich, C. E.; Frankowitz, S. H.; and Xintaras, C. Altered Function in Animals Inhaling Low Concentrations of Ozone and Nitrogen Dioxide. Am. Indust. Hyg. A. Quart. 25:246 (May-June), 1964.
70. Scheel, L. D.; Dobrogorski, O. J.; Mountain, J. T.; Svirbely, J. L.; and Stokinger, H. E. Physiologic, Biochemical, Immunologic and Pathologic Changes Following Ozone Exposure. J. Appl. Physiol. 14.1: 67 (Jan.), 1959.
71. McDonnell, H. B. Experiments with Ozone on Guinea Pigs. J. Assoc. Off. Agric. Chem. 13:19, 1930.
72. Brinkman, R.; Lamberts, H. B.; and Veninga, T. S. Radiometric Toxicity of Ozonized Air. Lancet 7325: 133 (Jan. 18), 1964.
73. Stokinger, H. E.; Wagner, W. D.; and Dobrogorski, O. J. Ozone Toxicity Studies. III. Chronic Injury to Lungs of Animals Following Exposure at a Low Level. Arch. Environ. Health 16:514, 1957.
74. Smith, L. E. Peroxyacetyl Nitrate Inhalation. Cardio-respiratory Effects. Ibid. 10:161 (Feb.), 1965.
75. Renzetti, N. A., and Schuck, E. A. Preliminary Observations on the Relationship Between Eye Irritation in Synthetic Systems and in the Atmosphere. J. Air Pollut. Contr. Ass. 11:191 (Mar.), 1961.
76. Renzetti, N. A. An Aerometric Survey of the Los Angeles Basin, August-November, 1954. Rep. No. 9, Air Pollution Foundation, Los Angeles, Calif. (July), 1955.
77. Coffin, D. L., and Blommer, E. J. Acute Toxicity of



- Irradiated Acute Exhaust. Its Indication by Enhancement of Mortality from Streptococcal Pneumonia. Arch. Environ. Health. In press.
78. Motley, H. L.; Smart, R. H.; and Leftwich, C. I. Effect of Polluted Los Angeles Air (Smog) on Lung

- Volume Measurements. J.A.M.A. 171,11:1469 (Nov.), 1959.
79. Bils, R. Ultrastructural Alterations of Alveolar Tissue of Mice. I. Due to Heavy Los Angeles Smog. Arch. Environ. Health 12:689 (June), 1966.

Mr. Jaffe is physical science administrator, National Center for Air Pollution Control, Public Health Service, Washington, D. C.

This paper was presented before the Occupational Health Section of the American Public Health Association at the Ninety-Fourth Annual Meeting in San Francisco, Calif., November 2, 1966.

## In England Now

A Running Commentary by Peripatetic Correspondents

Henry Sigerist, whose *Autobiographical Writings* has now appeared in an English edition, to the end of his life, never ceased to find inspiration from the sight of a blank page of his specially bound exercise books. His fountain-pen flowed, and the number of pages he set himself to write each day were covered with his characteristic widely spaced sprawling words. The lesson seems to be not to wait on inspiration. In this way one can perhaps ensure quantity. But quality? There lies the rub. Sigerist had no difficulty. To describe him as the "universal man" of the Renaissance is almost a cliché. For him there was no dichotomy between art and science. Indeed, if you wanted to sum up his lifework in a sentence, he placed medical history squarely in the framework of history as a whole. But history as a whole meant for him the everyday life of slaves, peasants, craftsmen, and labourers as well as the life of doctors, men of science, painters, musicians, architects, scholars, princes, and politicians. So that when he spoke about William Harvey his account of the contemporary scene seemed to make it inevitable that Harvey *had* to discover just as that time how the circulation worked. You might say that, like his fellow-countryman, the art historian Burckhardt, he contrived to make historical harmony out of seemingly unrelated and discordant historical figures and events.

When he came to our university in 1939 to give some guest lectures, people expected to be bored and instead were enchanted. His first lecture drew an audience of about twenty; his fifth had to be given in the largest hall and transmitted by loud-speaker to the next largest. I never saw him again. But contact with him sent me scurrying to books on the history of painting, music, and architecture—and to desultory lessons in Italian. I think he made me a better doctor; he certainly turned me into a better teacher.

(The Lancet, June 17, 1967, p. 1323.)